

Research papers relating to the core and the buffer of the Biosphere.

Allot, T.E.H, Harriman, R. and Battarbae, R.W. (1992). Reversibility of lake acidification at the Round Loch of Glenhead, Galloway, Scotland. <i>Environmental Pollution</i> Vol. 77, Issues 2–3, pp. 219–225.
Anderson, N.J. (1990). Variability of sediment diatom assemblages in an upland, wind-stressed lake (Loch Fleet, Galloway, S.W. Scotland). <i>Journal of Paleolimnology</i> Vol. 4, No. 1, pp. 43-59.
Armitage, H. (2010). Assessing the influence of environmental drivers on the current condition and recovery potential of <i>Racomitrium</i> heath. PhD thesis, University of Aberdeen.
Barnes, R.P., Anderson, T.B. and McCurry, J.A. (1987). Along-strike variation in the stratigraphical and structural profile of the Southern Uplands Central Belt in Galloway and Down. <i>Journal of the Geological Society</i> Vol. 144 No. 5, pp. 807-817.
Battarbee, R.W., Stevenson, A.C., Rippey, B., Fletcher, C., Natkanski, J., Wik, M. and Flower, R.J. (1989). Causes of Lake Acidification in Galloway, South-West Scotland: A Palaeoecological Evaluation of the Relative Roles of Atmospheric Contamination and Catchment Change for Two Acidified Sites with Non-Afforested Catchments. <i>Journal of Ecology</i> Vol. 77, No. 3, pp. 651-672.
Battarbee, R.W. (1989). Geographical research on acid rain. I. The acidification of Scottish lochs. <i>The Geographical Journal</i> Vol. 155, No.3, pp. 353 - 377.
Battarbee, R.W., Curtis, C.J. and Shilland, E.M. (2011). The Round Loch of Glenhead: Recovery from acidification, climate change monitoring and future threats. Scottish Natural Heritage Commissioned Report No. 469.
Battarbee, R.W., Flower, R.J., Stevenson, A.C. and Rippey, B. (1985). Lake acidification in Galloway: a palaeoecological test of competing hypotheses. <i>Nature</i> 314, pp. 350 - 352.
Birks, H.H. (1972). Studies in the Vegetational History of Scotland: II. Two Pollen Diagrams from the Galloway Hills, Kirkcudbrightshire. <i>Journal of Ecology</i> Vol. 60, No. 1, pp. 183-218.
Birse, E.L. and Robertson, J.S. (1976). Plant communities and soils of the lowlands and southern upland regions of Scotland. Macaulay Institute for Soil Research, Aberdeen, 87 pp.
Blyth, F.G.H. (1950). The sheared porphyrite dykes of South Galloway. <i>Quarterly Journal of the Geological Society</i> Vol. 105, pp. 393-423.
Boatman, D.J. (1983). The Silver Flowe National Nature Reserve, Galloway, Scotland. <i>Journal of Biogeography</i> Vol. 10, No. 3, pp. 163-275.
Boatman, D.J. and Tomlinson, R.W. (1973). The Silver Flowe I. Some structure and hydrological features of Brishie Bog and their bearing on pool formation. <i>Journal of Ecology</i> Vol. 61, pp.653-666.
Boatman, D.J. and Tomlinson, R.W. (1977). The Silver Flowe II. Features of the vegetation and stratigraphy of Brishie Bog and their bearing on pool formation. <i>Journal of Ecology</i> Vol. 65, pp. 531-546.
Boatman, D.J., Goode, D.A. and Hulme, P.D. (1981). The Silver Flowe III. Pattern development on Long Loch B and Craigeazel Mires. <i>Journal of Ecology</i> Vol. 69, pp.879-912.
Bray, R. (2008). Factors underlying differential ecosystem recovery from acidification of upland waters in the river Cree catchment area, Dumfries and Galloway, Scotland. PhD

thesis, University of Strathclyde.
Bullock, D.J. (1982). Aspects of the ecology of feral goats (<i>Capra</i> (domestic)) in the Southern Uplands. Doctoral thesis, Durham University.
Bullock, D.J., North, S. and Kerner, P. (1977). Census of the feral goat population on Cairnsmore of Fleet, New Galloway. Unpub. report to N. C. C., South-west Scotland Region, Balloch.
Bullock, D.J., North, S. and Williams, D. (1976). Feral goats on Cairnsmore of Fleet. Unpub. report to N. C. C., South-west Scotland Region, Balloch.
Burns, J.C., Coy, J.S., Tervet, D.J., Harriman, R., Morrison, B.R.S. and Quine, C.P. (1984). The Loch Dee Project: a study of the ecological effects of acid precipitation and forest management on an upland catchment in south-west Scotland. <i>Fisheries Management</i> Vol.15, pp. 145 - 167.
Calladine, J., Dott, H., Douglas, D. and Garner, G. (2009). Monitoring of moorland fringe biodiversity: the bird communities of the interface between conifer plantations and moorland in the Galloway Forest Park and their relationships with moorland fringe habitats. Scottish Natural Heritage Commissioned Report No.456.
Collen, P., Harriman, R., Morrison, B., Keay, E. and Watt, A. (2000). Restoration of a brown trout (<i>Salmo trutta</i> L.) population to Loch Enoch, an acidified Loch in Galloway, south-west Scotland. <i>Freshwater Forum</i> Vol.14, pp. 3-14.
Cook, D.R. and Weir, J.A. (1979). Structure of the Lower Palaeozoic rocks around Cairnsmore of Fleet, Galloway. <i>Scottish Journal of Geology</i> Vol.15, pp.187-202.
Cook, J.M., Edmunds, W.M. and Robins, N.S. (1991). Groundwater contribution to an acid upland lake (Loch Fleet, Scotland) and the possibilities for amelioration. <i>Journal of Hydrology</i> Vol. 125 No.1-2, pp. 111-128.
Cooper, D.M., Helliwell, R.C. and Coull, M.C. (2004). Predicting acid neutralizing capacity from landscape classification: application to Galloway, south-west Scotland. <i>Hydrological Processes</i> Vol. 18 No.3, pp. 455-471.
Dalziel, T.R.K., Howells, G., and Skeffington, R.A. (1992). The interaction of forest vegetation and soils with the aquatic environment; effects of catchment liming on lakes. <i>Studies in Environmental Science</i> Vol. 50, pp. 107-127.
Dalziel, T.R.K., Wilson, E.J. and Proctor, M.V. (1994). The effectiveness of catchment liming in restoring acid waters at Loch Fleet, Galloway, Scotland. <i>Forest Ecology and Management</i> Vol. 68 No. 1, pp.107-117.
Davies, E. J. M. (1982). Forestry in Dumfries and Galloway. <i>Forestry</i> Vol. 55 No.1, pp. 79-89.
Daysh, G.H.J. (1974). Geology. In: Galloway Forest Park. Forestry Commission Guide, Edlin, H.L. (Ed.), pp. 24-27. Her Majesty's Stationery Office, Edinburgh.
Dayton, N. (2007). Site Condition Monitoring of notified blanket bog features on Cairnsmore of Fleet SSSI, Dumfries and Galloway. Unpublished Report, Scottish Natural Heritage.
Dayton, N. and Rafferty, T. (2010). Site condition monitoring survey of upland and peatland notified features on designated sites in Dumfries and Galloway (Merrick Kells SAC and SSSI). A report to Scottish Natural Heritage by Quadrat Scotland. Contract 25029.

Dickson, R. C. (1994). Wintering Hen Harriers in West Galloway. Transactions of the Dumfriesshire and Galloway Natural History and Antiquarian Society Third Series Vol. 69, pp.3-6.
Dickson, R. C. (1995). Feeding rates of Hen Harriers in West Galloway. Scottish Birds Vol.18, pp. 104-106.
Dickson, R. C. (1996). The hunting behaviour of Merlins in winter in Galloway. Scottish Birds Vol. 18, pp. 165-169.
Dickson, R. C. (1999). Size of Merlin pellets in winter and summer in Galloway. Scottish Birds Vol. 20, pp. 31-33.
Dickson, R. C. (2003). Egg weight and growth of nestling Merlins in Dumfries & Galloway. British Birds Vol. 96, pp. 252-257.
Donoghue, D.N.M., Watt, P.J., Cox, N.J. and Wilson, J. (2007). Remote sensing of species mixtures in conifer plantations using LiDAR height and intensity data. Remote Sensing of Environment Vol. 110, No. 4, pp. 509–522.
Donoghue, D.N.M., Watt, P.J., Cox, N.J., Dunford, R.W., Wilson, J., Stables, S. and Smith, S. (2004). An evaluation of the use of satellite data for monitoring early development of young Sitka spruce plantation forest growth. Forestry Vol. 77 No.5, pp. 383-397.
Dowens, J.S. (2007). Fire damage assessment report: Silver Flowe NNR, Galloway Forest Park, Scotland. Report for Scottish Natural Heritage. School of Geosciences, University of Edinburgh.
Drakeford, T. Land use history of Cairnsmore of Fleet. Unpub. Report to N. C. C., South-west Scotland Region, Balloch.
Dunford, R. (2008). Forest impacts on freshwater acidification: an investigation of policy and practice in Galloway, S.W. Scotland. PhD thesis, Durham University.
Dunford, R.W., Donoghue, D.N.M. and Burt, T.P. (2012). Forest land cover continues to exacerbate freshwater acidification despite decline in sulphate emissions. Environmental Pollution Vol. 167, pp. 58–70.
Edlin, H.L., (1974). Galloway Forest Park. Forestry Commission Guide. Her Majesty's Stationary Office, Edinburgh, 79 pp.
Evers, S.L. (2008). The role of forest stream corridor characteristics in influencing stream and riparian ecology. PhD thesis, Glasgow University.
Farley, D.A. and Werritty, A. (1989). Hydrochemical budgets for the Loch Dee experimental catchments, south-west Scotland. Journal of Hydrology Vol. 109, pp. 351 - 368.
Ferguson, A., Prodöhl, P., Ade, R. (2012). Galloway & Carrick Trout Study 2010-2012. Native genetic diversity and anthropogenic influences on Galloway & Carrick brown trout (<i>Salmo trutta</i>) populations. Preliminary report.
Ferrier, R. C., Helliwell, R. C., Cosby, B. J., Jenkins, A. and Wright, R. F.(2001). Recovery from acidification of lochs in Galloway, south-west Scotland, UK: 1979-1998. Hydrology and Earth System Sciences Vol.5 No. 3, pp. 421-432.

<p>Ferrier, R.C., Jenkins, A., Cosby, B.J., Helliwell, R.C., Wright, R.F. (1995) Land use and deposition interactions: an evaluation of the effects of future N deposition scenarios on the Galloway Region of south-west Scotland using coupled sulphur and nitrogen model (MAGIC-WAND). <i>Water, Air and Soil Pollution</i>, Vol. 85 No.2, pp. 707 - 712.</p>
<p>Flower, R.J. (1986). The relationship between surface sediment diatom assemblages and pH in 33 Galloway lakes: Some regression models for reconstructing pH and their application to sediment cores. <i>Hydrobiologia</i> Vol. 143, No 1, pp.93-103.</p>
<p>Flower, R.J. and Battarbee, R.W. (1983). Acid Lakes in the Galloway Uplands, South West Scotland: catchments, water quality and sediment characteristics. (PRU Research Report 6). Palaeoecology Research Unit London.</p>
<p>Flower, R.J. and Battarbee, R.W. (1985). The morphology and biostratigraphy of <i>Tabellaria quadrisepitata</i> (Bacillariophyceae) in acid waters and lake sediments in Galloway, Southwest Scotland. <i>British Phycological Journal</i> Vol. 20 No.1, pp. 69-80.</p>
<p>Flower, R.J., Battarbee, R.W. and Appleby, P.G. (1987). The recent palaeolimnology of acid lakes in Galloway, south-west Scotland: diatom analysis, pH trends, and the role of afforestation. <i>Journal of Ecology</i> Vol. 75, pp. 797-824.</p>
<p>Flower, R.J., Rippey, B. and Tervet, D.J. (1986). Thirty four Galloway Lochs: bathymetry, water quality and surface sediment diatom assemblages. (PRU Research Report 14). Palaeoecology Research Unit London.</p>
<p>Forestry Commission, (1999). Inventory report. National Inventory of Woodland and Trees. Scotland-Dumfries and Galloway region. Part 1 - Woodlands of two hectares and over. Woodlands Survey, Forest Research, pp. 5–12.</p>
<p>Gale, M.F. and Anderson, A.B. (1994). High elevation planting in Galloway. <i>Scottish Forestry</i> Vol. 38 No.1, pp. 3-17.</p>
<p>Gardiner, C.I. and Reynolds, S.H. (1932). The Loch Doon 'Granite' Area, Galloway. <i>Quarterly Journal of the Geological Society</i>, Vol. 88, 1-NP.</p>
<p>Ginns, M., Dinsmore, G. (2010). Riparian mammals on the upper Cree catchment. Research report. Lutra Ecological Services.</p>
<p>Grieve, I. C. (1990). Variations in the chemical composition of the soil solution over a four year period at an upland site in s.w. Scotland. <i>Geoderma</i>, Vol. 46, pp. 351 - 362.</p>
<p>Grieve, I.C. (1990). Effects of catchment liming and afforestation on the concentration and fractional composition of aluminium in the Loch Fleet catchment, SW Scotland. <i>Journal of Hydrology</i> Vol. 115 No. 1–4, pp. 385–396.</p>
<p>Harriman, R., Morrison, B.R.S., Birks, H.J.B., Christie, A.E.G., Collen, P. and Watt, A.W. (1995). Long term chemical and biological trends in Scottish streams and lochs. <i>Water, Air and Soil Pollution</i> Vol. 85, pp. 701-706.</p>
<p>Harriman, R., Morrison, B.R.S., Caines, L.A., Collen, P. and Watt, A.W. (1987). Long-term changes in fish populations of acid streams and lochs in Galloway South West Scotland. <i>Water, Air, & Soil Pollution</i> Vol. 32 No 1-2, pp. 89-112.</p>
<p>Harriman, R., Watt, A.W., Christie, A.E.G., Collen, P., Moore, D.W., McCartney, A.G., Taylor, E.M. and Watson, J. (2001). Interpretation of trends in acidic deposition and surface water chemistry in Scotland during the past three decades. <i>Earth System Sciences</i> Vol. 5 No.3, pp. 407–420.</p>

Helliwell, R.C., Ferrier, R.C., Johnston, L., Goodwin, J. and Doughty, R. (2001). Land use influences on acidification and recovery of freshwaters in Galloway, south-west Scotland. <i>Hydrology and Earth System Sciences</i> Vol. 5 No. 3, pp. 451-458.
Helliwell, R.C., Ferrier, R.C. and Kernan, M.R. (2001). Interaction of nitrogen deposition and land use on soil and water quality in Scotland: issues of spatial variability and scale. <i>Science of The Total Environment</i> Vol. 265 No. 1-3, pp. 51-63.
Henderson, D.J., Malcolm, A., Nolan, A.J., Hewison, R.L., McLeod, J. and Alexander, J. (2004). Habitat Condition Assessment and Modelling for Appropriate Grazing Management of Merrick-Kells SSSI, and Heather Condition Assessment for Cairnsmore of Fleet NNR. A report for Scottish Natural Heritage. Macaulay Research Consultancy Services.
Hewison, R.L., Henderson, D.J., Malcolm, A., Nolan, A.J. and Alexander, J. (2006). Habitat condition assessment of Merrick Kells (Site of Special Scientific Interest and Candidate Special Area of Conservation) and Cairnsmore of Fleet (Site of Scientific Interest and National Nature Reserve). Scottish Natural Heritage Commissioned Report No. 196 (ROAME No. F02LK12).
Holland, J.P., Pollock, M., Waterhouse, T., Morgan-Davies, C., Bibby, H., Stewart, S. and Armstrong, H.M. (2010). Developing guidance for managing extensive upland grazing where habitats have differing requirements. Scottish Natural Heritage Commissioned Report No.402.
Horsfield, D. (2007). Fire damage assessment report May 2007: Merrick Kells Special Area of Conservation. Report for Scottish Natural Heritage.
Howells, G. and Dalziel, T. (1995). A decade of studies at Loch Fleet, Galloway (Scotland): A catchment liming project and restoration of a brown trout fishery. <i>Freshwater Forum</i> Vol. 5 No.1, pp. 4-38.
Howells, G., Dalziel, T.R.K. and Turnpenny, A.W.H. (1992). Loch Fleet: liming to restore a brown trout fishery. <i>Environmental Pollution</i> , pp.131-139.
J. Watson (1979). Food of Merlins nesting in young conifer forest. <i>Bird Study</i> , Vol. 26 No. 4, pp. 253-258.
Jardine, W.G. (1959). River development in Galloway. <i>Scottish Geographical Magazine</i> Vol. 75 No. 2, pp. 65-75.
Jardine, W.G. (1967), Landscape evolution in Galloway. <i>Trans. J. Proc. Dumfries. Galloway Natural History and Antiquarian Society</i> .
Jeffries, G. (1971). Goat Park. Unpub. report to Forestry Commission, Newton Stewart.
Jones, V.J, Stevenson, A.C., and Battarbee, R.W. (1989). Acidification of Lakes in Galloway, South West Scotland: A Diatom and Pollen Study of the Post-Glacial History of the Round Loch of Glenhead. <i>Journal of Ecology</i> Vol. 77, No. 1, pp. 1-24.
Langan, S.J. and Hirst, D. (2004). An analysis of the long-term variation in stream water quality for three upland catchments at Loch Dee (Galloway, S.W. Scotland) under contrasting land management. <i>Hydrology and Earth System Sciences</i> Vol. 8, No. 3, pp.422-435.
Langan, S.J., (1987). Episodic acidification of streams at Loch Dee, S.W.Scotland. <i>Transactions of the Royal Society of Edinburgh</i> Vol. 78, pp. 393-398.

Leech, A.F. And Nisbet, T.R. (1989). Changes in rain and soil water chemistry on passage through forested and unforested site at Loch fleet, Galloway.
Lees, F.M. (1992). Water chemistry and hydrology of Loch Dee. Occasional publication, Solway River Purification Board, Dumfries, Scotland.
Lees, F.M. (1995). Have emission reductions led to improvements in water quality at Loch Dee in Galloway? <i>Water, Air, & Soil Pollution</i> Vol. 85 No. 2, pp. 695-700.
McCormick, A. (1932). Galloway. The spell of its hills and glens. John Smith, Glasgow.
McIntosh, R. (1978). Response of Sitka spruce to remedial fertilisation in Galloway [Scotland]. <i>Scottish Forestry</i> Vol.32 No.4 pp. 271-283.
M'Donald, J. (1927). The tailless trout of Loch Enoch. <i>Dumfries and Galloway Natural History and Antiquarian Society</i> XIV, pp. 299-309.
Mearns, R. (2001). Juniper, <i>Juniperus communis</i> in Dumfries and Galloway. In the <i>Transactions of the Dumfriesshire and Galloway Natural History and Antiquarian Society</i> , LXXV.
Miller, K.F., Quine, C.P. and Hunt, J. (1987). The Assessment of Wind Exposure for Forestry in Upland Britain. <i>Forestry</i> Vol. 60 No. 2, pp. 179-193.
Milne, J.A., Birch, C.P.D., Hester, A.J., Armstrong, H.M. and Robertson, A. (1998). The impact of vertebrate herbivores on the natural heritage of the Scottish Uplands - a review. <i>Scottish Natural Heritage Review</i> No 95.
Mohamed, A. and Malthus, T. (2007). Characterising peatland surface heterogeneity using hyperspectral airborne remote sensing data. <i>Proceedings 5th EARSeL Workshop on Imaging Spectroscopy</i> . Bruges, Belgium, April 23 - 25 2007.
Morrison, B.R.S. (1997). Stomach contents of trout, <i>Salmo trutta</i> L., caught by angling on Loch Dee, an acid water in south west Scotland. <i>Fisheries Management and Ecology</i> Vol. 4, pp. 217-222.
Nature Conservancy Council (1975). The vegetation of Cairnsmore of Fleet. Unpub. report by V. Meek.
Orchel, J. (1992). Forest Merlins in Scotland: their requirements and management: a report on the Galloway Forests Merlin Project (1986-89). The Hawk and Owl Trust, London.
Petty, S. J. and Anderson, D. (1986). Breeding by Hen Harriers <i>Circus cyaneus</i> on restocked sites in upland forests, <i>Bird Study</i> , Vol. 33 No. 3, pp.177-179.
Piper, J. D. A., McArdle, N. J. and Almaskeri, Y. (2007). Palaeomagnetic study of the Cairnsmore of Fleet Granite and Criffel-Dalbeattie granodiorite contact aureoles: Caledonian tectonics of the Southern Uplands of Scotland and Devonian palaeogeography. <i>Geological Magazine</i> Vol. 144 No.5, pp. 811-835.
PlaceFirst (2011). Agri-tourism in Southern Scotland. Scottish Natural Heritage Commissioned Report No. 463.
Puhr, C.B. (1997). Catchment afforestation, surface water acidification, and salmonid populations in Galloway, South West Scotland. PhD thesis, Durham University.
Puhr, C.B. and Donoghue, D.N.M. (2000). Remote sensing of upland conifer plantations using Landsat TM data: A case study from Galloway, south-west Scotland. <i>International Journal of Remote Sensing</i> Vol. 21, Issue 4, pp. 633-647.

Puhr, C.B., Donoghue, D.N.M., Stephen, A.B., Tervet, D.J. and Sinclair, C. (2000). Regional Patterns of Streamwater Acidity and Catchment Afforestation in Galloway, SW Scotland. <i>Water, Air, & Soil Pollution</i> Vol.120 No.1-2, pp. 47-70.
Quinn, C.H., Fraser, E.D.G., Hubacek, K. and Reed, M.S. (2010). Property rights in UK uplands and the implications for policy and management. <i>Ecological Economics</i> Vol. 69, Issue 6, pp. 1355–1364.
Ratcliffe, D.A. (2007). <i>Galloway and the Borders</i> . New Naturalists, Harper Collins, London.
Ratcliffe, D.A. and Walker, D. (1958). The Silver Flowe, Galloway, Scotland. <i>Journal of Ecology</i> Vol. 46, No. 2, pp. 407-445.
Raven, P.J. (1988). Occurrence of sphagnum moss in the sublittoral of several small oligotrophic lakes in Galloway, Southwest Scotland. <i>Aquatic Botany</i> Vol. 30 No. 3, pp. 223–230.
Rees, R.M. and Ribbens, J.C. H. (1995). Relationships between afforestation, water chemistry and fish stocks in an upland catchment in south west Scotland. <i>Water, Air, & Soil Pollution</i> Vol. 85 No. 2, pp.303-308.
Ribbens, J.C.H. and Graham, J.L. (2004). Strategy for the containment and possible eradication of American signal crayfish (<i>Pacifastacus leniusculus</i>) in the River Dee catchment and Skyre Burn catchment, Dumfries and Galloway. Scottish Natural Heritage Commissioned Report No. 014. (ROAME No. F02LK05).
Riddle, G., and Sheppard., G. (1999). Large kestrel clutch sizes in Southwest Scotland 1997-98. <i>Scottish Birds</i> Vol. 20:43.
Shaw, G. (1995). Habitat selection by short-eared owls <i>Asio flammeus</i> in young coniferous forests. <i>Bird Study</i> , Vol.42 No.2, pp.158-165.
Shaw, G. and Livingstone, J. (1994). The pine marten–its reintroduction and subsequent history in the Galloway Forest Park. <i>Transactions of the Dumfries and Galloway Natural History and Antiquarian Society</i> .
Smith, M. A. (1990). The Ecophysiology of Epilithic Diatom Communities of Acid Lakes in Galloway, Southwest Scotland <i>Phil. Transactions of the Royal Society of London</i> Vol. 327 No.1240, pp. 251-257.
SNH (2002). <i>Western Southern Uplands and Inner Solway</i> . Natural Heritage Futures series. Scottish Natural Heritage, Battleby.
Stone, P. (ed.) (1996). <i>Geology in south-west Scotland: An excursion guide</i> . British Geological Survey.
Tervet, D.J. and Harriman, R. (1988). Changes in pH and calcium after selective liming in the catchment of Loch Dee, a sensitive and rapid turnover loch in south-west Scotland. <i>Aquaculture and Fisheries Management</i> Vol. 19, pp. 191 - 203.
Theaker, J. (1979). Census of the feral goat population on Cairnsmore of Fleet. Unpub. report to N. C. C., South-west Scotland Region, Balloch.
Theaker, J. (1980). Census of the feral goat population, Cairnsmore Fleet N. N. R. Unpub. report to N. C. C., South-west Scotland Region, Balloch.
Turnpenny, A.W.H., Fleming, J.M. and Wood, R. (1995). The brown trout population at Loch Fleet eight years after liming. <i>Chemistry and Ecology</i> Vol. 9, pp. 179-191.

Warren, C. (2000). 'Birds, bogs and forestry' revisited: The significance of the flow country controversy. *Scottish Geographical Journal* Vol. 116 No. 4, pp. 315-337.

Willet, J. and Corcoran, S. (2010). Site Condition Monitoring of dragonflies on selected Site of Special Scientific Interest (SSSIs). Preliminary Survey Report for Blackwood of Rannoch, Glen Strathfarrar, Merrick Kells, River Dee, Kenmure Holms and Sunart SSSIs.

Williams, T.M. (1991). A sedimentary record of the deposition of heavy metals and magnetic oxides in the Loch Dee basin, Galloway, Scotland, since c. AD 1500. *The Holocene* Vol. 1 No. 2, pp.142-151.

Williams, T.M. (1988). The geochemistry and magnetic mineralogy of Holocene strata in acid basin: Case-studies of granitic catchments in the Galloway and Rannochmoor regions of Scotland. PhD thesis, University of Edinburgh.

Williamson, K. (1976). Bird-life in the wood of Cree, Galloway [Scotland]. *Quarterly Journal of Forestry* Vol. 70 No.4, pp. 206-216.

Willis, K.G. and Benson, J.F. (1989). Recreational Values of Forests. *Forestry* Vol.62 No.2, pp. 93-111.

Wright, R.F. and Henriksen, A. (1980). Regional survey of lakes and streams in southwestern Scotland, April 1979. Internal Report IR 72/80, SNSF-Project, Ås, Norway, 63 pp.

Wright, R.F., Cosby, B.J., Ferrier, R.C., Jenkins, A., Bulger, A.J. and Harriman, R. (1994). Changes in acidification of lochs in Galloway, southwestern Scotland, 1979–1988: The MAGIC model used to evaluate the role of afforestation, calculate critical loads, and predict fish status. *Journal of Hydrology* Vol. 161, Issues 1–4, pp. 257–285.

Yang, H. (2009). Spatial and temporal patterns of benthic diatoms in an upland Scottish loch (the Round Loch of Glenhead): with reference to recent environmental changes. 370pp. PhD thesis, Department of Geography. University College London, London.

